IS MY CHILD TOO ILL FOR SCHOOL?

yes until...

CHICKENPOX	At least 5 days from the onset of the rash and until all blisters have crusted over.
DIARRHOEA AND VOMITING	48 hours after their last episode.
HIGH TEMPERATURE	If your child has a high temperature, keep them off school until it goes away.
IMPETIGO	Their sores have crusted and healed, or 48 hours after they started antibiotics.
MEASLES	4 days after the rash first appeared.
MUMPS	5 days after their swelling started.
SCABIES	They've had their first treatment.
SCARLET FEVER	24 hours after they started taking antibiotics.
WHOOPING COUGH	48 hours after they started taking antibiotics.



NO



CONJUNCTIVITIS	You don't need to keep your child away from school if they have conjunctivitis. Do get advice from your pharmacist. Encourage your child not to rub their eyes and to wash their hands regularly.
COUGHS & COLDS	It's fine to send your child to school with a minor cough or common cold.
HAND, FOOT & MOUTH	If they seem well enough to go to school, there's no need to keep them off. Encourage your child to throw away any used tissues straight away and to wash their hands regularly.
HEAD LICE	There's no need to keep your child off school. You can treat head lice and nits without seeing your GP.
THREADWORMS	Speak to your pharmacist, who can recommend treatment.
SORE THROAT TONSILLITIS	You can still send your child to school if they have a sore throat. But if they also have a high temperature, they should stay at home until it goes away.
SLAPPED CHEEK	You don't need to keep your child off school if they have slapped cheek syndrome because, once the rash appears, they're no longer infectious.